

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 February 2001 (08.02.2001)

PCT

(10) International Publication Number
WO 01/09114 A1

(51) International Patent Classification⁷: **C07D 325/00**,
C12P 17/18, A61K 31/335

Fernando [ES/ES]; Josefa Valcarcel; 3, E-28027 Madrid (ES).

(21) International Application Number: PCT/US00/20918

(74) Common Representative: **MERCK & CO., INC.**; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(22) International Filing Date: 31 July 2000 (31.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/146,994 3 August 1999 (03.08.1999) US

(71) Applicant (for all designated States except US): **MERCK & CO., INC.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **DOMBROWSKI, Anne** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **POLISHOOK, Jon, D.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **SINGH, Sheo** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **ZINK, Deborah, L.** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **HAZUDA, Daria** [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065-0907 (US). **TERAN, Ana** [ES/ES]; Josefa Valcarcel, 3, E-28027 Madrid (ES). **PELAEZ,**

Published:

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/09114 A1

(54) Title: HIV INTEGRASE INHIBITORS

(57) Abstract: Novel tetracyclic aromatic ketones are natural product compounds useful in the inhibition of HIV integrase, the prevention or treatment of infection by HIV and the treatment of AIDS, either as compounds, pharmaceutically acceptable salts, pharmaceutical composition ingredients, whether or not in combination with other antivirals, immunomodulators, antibiotics or vaccines. Methods of treating AIDS and methods of preventing or treating infection by HIV are also described. Further, the novel fungal cultures MF6388 (ATCC 74478), *Sterile Fungus*, and MF6591 (ATCC 74477), *Ascochyta sp.* are also disclosed. Further, the cultures *Sterile Fungus* MF6388 (ATCC 74478) and *Ascochyta sp.* MF6591 (ATCC 74477) are also disclosed, as well as processes for making compounds of the present invention employing the cultures.

TITLE OF THE INVENTION
HIV INTEGRASE INHIBITORS

BACKGROUND OF THE INVENTION

5 A retrovirus designated human immunodeficiency virus (HIV) is the etiological agent of the complex disease that includes progressive destruction of the immune system (acquired immune deficiency syndrome; AIDS) and degeneration of the central and peripheral nervous system. This virus was previously known as LAV, HTLV-III, or ARV. A common feature of retrovirus
10 replication is the insertion by virally-encoded integrase of proviral DNA into the host cell genome, a required step in HIV replication in human T-lymphoid and monocytoïd cells. Integration is believed to be mediated by integrase in three steps: assembly of a stable nucleoprotein complex with viral DNA sequences; cleavage of two nucleotides from the 3' termini of the linear proviral DNA;
15 covalent joining of the recessed 3' OH termini of the proviral DNA at a staggered cut made at the host target site. The fourth step in the process, repair synthesis of the resultant gap, may be accomplished by cellular enzymes.

 Nucleotide sequencing of HIV shows the presence of a pol gene in one open reading frame [Ratner, L. et al., Nature, 313, 277(1985)]. Amino acid
20 sequence homology provides evidence that the pol sequence encodes reverse transcriptase, integrase and an HIV protease [Toh, H. et al., EMBO J. 4, 1267 (1985); Power, M.D. et al., Science, 231, 1567 (1986); Pearl, L.H. et al., Nature, 329, 351 (1987)]. All three enzymes have been shown to be essential for the replication of HIV.

25 It is known that some antiviral compounds which act as inhibitors of HIV replication are effective agents in the treatment of AIDS and similar diseases, e.g., azidothymidine or AZT. Applicants demonstrate that the compounds of this invention are inhibitors of HIV integrase. The applicants additionally demonstrate that inhibition of integrase in vitro is a direct result of
30 inhibiting the strand transfer reaction catalyzed by the recombinant integrase in vitro. The particular advantage of the present invention is highly specific inhibition of HIV integrase and HIV replication. The compounds of the present invention inhibit integrases of closely related lentiviruses such as HIV 2 and SIV, but not integrases from more distantly related retroviruses, for example RSV.
35 These compounds do not inhibit binding or catalysis of other nucleic acid binding

proteins, including enzymatic reactions such as those catalyzed by HIV reverse transcriptase, HIV RNase H, Influenza transcriptase, Hepatitis C polymerase, Yeast DNA polymerase, DNase I, Eco RI endonuclease, or mammalian polymerase II.

5 Zhao et al., (J. Med Chem. vol. 40, pp. 937-941 and 1186-1194 (1997)) describe hydrazide and arylamide HIV integrase inhibitors. Bis-catechols useful for inhibiting HIV integrase are described in LaFemina et al. (Antimicrobial Agents & Chemotherapy, vol. 39, no. 2, pp. 320-324, February 1995).

10 Applicants have discovered that certain tetracyclic aromatic ketones are potent inhibitors of HIV integrase. These compounds are useful for the treatment of AIDS or HIV infections.

BRIEF DESCRIPTION OF THE INVENTION

15 Compounds of Formula I, as herein defined, are disclosed. These compounds are useful in the inhibition of HIV integrase, the prevention of infection by HIV, the treatment of infection by HIV and in the treatment of AIDS and/or ARC, either as compounds, pharmaceutically acceptable salts or hydrates (when appropriate), pharmaceutical composition ingredients, whether or not in combination with other antivirals, anti-infectives, immunomodulators, antibiotics or vaccines.


20 Methods of treating AIDS, methods of preventing infection by HIV, and methods of treating infection by HIV are also disclosed. Further, the cultures *Sterile Fungus* MF6388 (ATCC 74478) and *Ascochyta sp.* MF6591 (ATCC 74477) are also disclosed, as well as processes for making compounds of the present invention employing the cultures.

25

DETAILED DESCRIPTION OF THE INVENTION

This invention is concerned with compounds of Formula I, combinations thereof, or pharmaceutically acceptable salts thereof, in the inhibition of HIV integrase, the prevention or treatment of infection by HIV and in the treatment of

30 the resulting acquired immune deficiency syndrome (AIDS). Compounds of Formula I are defined as follows:



(I)

wherein:

R is selected from:


5

(a) $-\text{CH}_2\text{OH}$, and

(b) $-\text{CH}(=\text{O})$;


or a pharmaceutically acceptable salt thereof.

Particular compounds of structural Formula I include:



10

Compound A, and



Compound B.

Also included within the present invention are pharmaceutical compositions useful for inhibiting HIV integrase, comprising an effective amount of a compound of this invention, and a pharmaceutically acceptable carrier.

Pharmaceutical compositions useful for treating infection by HIV, or for treating

15

AIDS or ARC, are also encompassed by the present invention, as well as a method of inhibiting HIV integrase, and a method of treating infection by HIV, or of treating AIDS or ARC. Additionally, the present invention is directed to a pharmaceutical composition comprising a therapeutically effective amount of a compound of the present invention in combination with a therapeutically effective amount of an AIDS treatment agent selected from:

- (1) an AIDS antiviral agent,
- (2) an anti-infective agent, and
- (3) an immunomodulator.

The compounds of the present invention may have asymmetric centers and may occur, except when specifically noted, as mixtures of stereoisomers or as individual diastereomers, or enantiomers, with all isomeric forms being included in the present invention.

This invention also discloses the culture MF6388 (ATCC 74478) identified as *Sterile Fungus*. A culture of MF6388 (ATCC 74478) is defined as substantially free of its natural soil contaminants and capable of forming a compound of Formula (I) in a recoverable amount. The culture should be free from viable contaminating microorganisms deleterious to the production of a compound of Formula (I). A biologically pure culture of MF6388 (ATCC 74478) may also be employed. In one embodiment, the present invention includes a culture of MF6388 (optionally biologically pure), or a mutant thereof, capable of producing in a recoverable amount a compound of Formula (I).

Suitable mutant strains of MF6388 can be obtained by chemically induced mutagenesis using mutagens such as nitrosoguanidine, 1-methyl-3-nitro-1-nitrosoguanidine, ethyl methane sulfonate, 2-aminopurine, and the like. Mutant strains can also be obtained by radiation-induced mutagenesis, such as by irradiation with ultraviolet light (e.g., using a germicidal lamp), X-rays, or gamma rays (e.g., using a cobalt-60 source). Recombinant DNA techniques such as protoplast fusion, plasmid incorporation, gene transfer and the like may also be employed. Further description of mutagenic techniques can be found in Vinci and Bing, "Strain Improvement by Nonrecombinant Methods", in *Manual of Industrial Microbiology and Biotechnology* 1999, 2d edition, edited by Demain et al., ASM Press, 103- 113; and in Carlton and Brown, "Gene Mutation", Chapter 13 in *Manual of Methods for General Bacteriology* 1985, edited by Gerhardt et al., ASM Press, 222-229.

In addition, compounds of the present invention may be prepared by fermentation of the culture MF6388, ATCC 74478.

The present invention also relates to the preparation of compounds of structural Formula I comprising:

- 5 (a) fermenting a culture of MF6388 (ATCC 74478), *Sterile Fungus* or a mutant thereof to produce a fermentation broth,
- (b) extracting the fermentation broth with an organic solvent,
- (c) isolating the compounds of structural Formula I.

The compounds of structural Formula I are preferably isolated by partitioning the fermentation extract between the organic solvent and water, followed by size
10 exclusion chromatography and normal or reverse-phase chromatography.

This invention also discloses the culture MF6591 (ATCC 74477) identified as *Ascochyta sp.* A culture of MF6591 (ATCC 74477) is defined as substantially free of its natural soil contaminants and capable of forming a compound
15 of Formula (I) in a recoverable amount. The culture should be free from viable contaminating microorganisms deleterious to the production of a compound of Formula (I). A biologically pure culture of MF6591 (ATCC 74477) may also be employed. In one embodiment, the present invention includes a culture of MF6591 (optionally biologically pure), or a mutant thereof, capable of producing in a
20 recoverable amount a compound of Formula (I). Suitable mutant strains of MF6591 can be obtained using the methods disclosed above for obtaining mutants of MF6388.

In addition, compounds of the present invention may be prepared by fermentation of the culture MF6591, ATCC 74477

The present invention also relates to the preparation of compounds of structural Formula I comprising:
25

- (a) fermenting a culture of MF6591 (ATCC 74477), *Ascochyta sp.* or a mutant thereof to produce a fermentation broth,
- (b) extracting the fermentation broth with an organic solvent,
- (c) isolating the compounds of structural Formula I.

30 The compounds of structural Formula I are preferably isolated by partitioning the fermentation extract between the organic solvent and water, followed by size exclusion chromatography and normal or reverse-phase chromatography.

When any variable (e.g., R¹, R², etc.) occurs more than one time in any constituent or in Formula I, its definition on each occurrence is independent of its

definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

The compounds of the present inventions are useful in the inhibition of HIV integrase, the prevention or treatment of infection by human immunodeficiency virus (HIV) and the treatment of consequent pathological conditions such as AIDS. Treating AIDS or preventing or treating infection by HIV is defined as including, but not limited to, treating a wide range of states of HIV infection: AIDS, ARC (AIDS related complex), both symptomatic and asymptomatic, and actual or potential exposure to HIV. For example, the compounds of this invention are useful in treating infection by HIV after suspected past exposure to HIV by e.g., blood transfusion, exchange of body fluids, bites, accidental needle stick, or exposure to patient blood during surgery.

The compounds of this invention are useful in the preparation and execution of screening assays for antiviral compounds. For example, the compounds of this invention are useful for isolating enzyme mutants, which are excellent screening tools for more powerful antiviral compounds. Furthermore, the compounds of this invention are useful in establishing or determining the binding site of other antivirals to HIV integrase, e.g., by competitive inhibition. Thus, the compounds of this invention are commercial products to be sold for these purposes.

The present invention also provides for the use of a compound of structural Formula (I) to make a pharmaceutical composition useful for inhibiting HIV integrase and in the treatment of AIDS or ARC.

Applicants have discovered that compounds of structural Formula (I), are useful for inhibiting HIV integrase.

ATCC Deposit of MF6388 (ATCC 74478), identified as *Sterile Fungus*.

Before the U.S. filing date of the present application, a sample of MF6388 (ATCC 74478), *Sterile Fungus*, was deposited at the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209, United States of America, under the terms of the Budapest Treaty. The culture access designation is 74478. This deposit will be maintained in the ATCC for at least 30 years and will be made available to the public upon the grant of a patent disclosing it. It should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by government action.

ATCC Deposit of MF6591 (ATCC 74477), identified as *Ascochyta sp.*

Before the U.S. filing date of the present application, a sample of MF6388 (ATCC 74477), *Ascochyta sp.*, was deposited at the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, VA 20110-2209, United States of America, under the terms of the Budapest Treaty. The culture access designation is 74477. This deposit will be maintained in the ATCC for at least 30 years and will be made available to the public upon the grant of a patent disclosing it. It should be understood that the availability of a deposit does not constitute a license to practice the subject invention in derogation of patent rights granted by government action.

10 General Characteristics and Descriptions of MF6388 (ATCC 74478), *Sterile Fungus* and MF6591 (ATCC 74477), *Ascochyta sp.*

In the following descriptions, MF6388 and MF6591 were edge inoculated with a 5 mm diameter plug on 2,100 mm petri dishes for each the following growth media. All cultures were incubated for 20 days at 25°C and 67% relative humidity in 12 hr photoperiod in fluorescent light unless otherwise indicated. In addition, all capitalized color names are from Ridgway, Color Standards and Nomenclature, (Published by author, Washington D.C., 1912) 43p. + 53.

Sterile Fungus (MF6388)

MF6388 was isolated from herbivore dung collected in New Mexico.
20 On oatmeal agar (Difco) culture mat attaining a diameter of 40 mm. Culture mat woolly from inoculation point to midway through the colony then thickly cottony until margin, grey-green (Pale Olive-Gray, Olive Gray), margin entire, white. Exudate forming as large, clear to light brown droplets near inoculation point and smaller, clear droplets closer to margin area. Reverse dark brown (Drab, Hair
25 Brown). Soluble pigment absent.

On potato-dextrose agar (Difco), culture mat attaining a diameter of 64 mm. Culture mat densely cottony, more dense at the inoculation point than closer to the margin, white aerial mycelium over green (Sage Green, Slate Olive) mycelium, margin entire, white. Exudate forming as sparse, clear droplets. Soluble pigment
30 dark red brown (Dark Indian Red, Indian Red).

On cornmeal agar (Difco), culture mat attaining a diameter of 75 mm (plate covered). Colony mat appressed, uncolored to slightly white. Margin entire, hyaline. Exudate, reverse and soluble pigment absent.

On YME agar (malt extract, 10.0 g; yeast extract, 4.0 g; dextrose, 4.0
35 g; agar, 20.0 g, distilled water, 1L) culture mat attaining a diameter of 66 mm.

Culture mat woolly, forming dense tufts, tufts becoming smaller towards margin area, white throughout, margin entire, white. Exudate forming as medium sized, light brown to sparse (at margin), clear droplets. Soluble pigment dark red brown (Dark Indian Red, Indian Red). No growth at 37°C, in the dark and no humidity control.

- 5 Microscopic: Hyphae hyaline, branched, thin-walled, usually 2 - 4 µm wide. No sporulation observed.

Ascochyta sp. (MF6591)

MF6591 was isolated from *Urtica urens* collected in Ontigola, Aranjuez, Spain.

- 10 On oatmeal agar (Difco) colony mat attaining a diameter of 40 mm. Culture mat cottony, dissected, light gray (Pale Neutral Gray, Gull Gray), abundant black fruiting bodies oozing light pink near inoculation point. Margin entire, light brown. Reverse dark gray (Deep Purplish Gray, Purplish Gray). Soluble pigment and exudate absent.

- 15 On potato-dextrose agar (Difco), colony mat attaining a diameter of 33 mm. Culture mat cottony, consistent throughout, gray-green (Olive Gray, Light Olive-Gray, Deep Olive-Gray), sparse white aerial mycelium near inoculation point. Soluble pigment deep red-brown (Indian Red, Deep Indian Red). Reverse and exudate absent.

- 20 On cornmeal agar (Difco), colony mat attaining a diameter of 45 mm. Culture mat appressed, uncolored. Fruiting bodies black, clustered around inoculation plug in contact with agar, immature fruiting bodies scattered throughout culture mat, abundant near inoculation point. Reverse, soluble pigment and exudate absent.

- 25 On YME agar (malt extract, 10.0 g; yeast extract, 4.0 g; dextrose, 4.0 g; agar, 20.0 g, distilled water, 1L) attaining a diameter of 33 mm. Culture mat cottony to thick velvety, light orange to pink (Vinaceous Pink, Shell Pink), sections near inoculation point collapsed, blackish. Margin region very faint yellow-brown (Light Buff, Pale Ochraceous Buff), very edge of margin entire, uncolored, submerged. Soluble pigment dark red-brown to light red-brown (Chestnut Brown, Cinnamon-Rufous). Exudate and reverse absent. At 37°C, in the dark and no humidity control, no growth.

- 30 Microscopic: Hyphae hyaline to light brown, thin-walled, branched, septate, 4 - 5 µm wide. Conidiomata pycnidial, thin-walled, black, spherical, mostly immersed in agar, 150 - 200 µm, mature conidia oozes as a pink mass from ostiole.

Conidiogenous cells phialidic, 1-celled, hyaline, 5 - 10 x 4 - 5 μ m. Conidia, hyaline, cylindrical, 1 medially septate, thin-walled, 10 - 15 x 3 - 4 μ m.

In general, MF6388 (ATCC 74478) is a strain cultured on a solid medium, or in an aqueous nutrient medium containing sources of assimilable carbon and nitrogen. For example, the cultures may be grown under submerged aerobic conditions (e.g., shaking culture, submerged culture, etc.)

In general, MF6591 (ATCC 74477) is strain that may be cultured on a solid medium, or in an aqueous nutrient medium containing sources of assimilable carbon and nitrogen. For example, the cultures may be grown under submerged aerobic conditions (e.g., shaking culture, submerged culture, etc.)

The sources of carbon in the nutrient medium are carbohydrates such as glucose, xylose, galactose, glycerin, starch, sucrose, dextrin, and the like. Other sources which may be included are maltose, rhamnose, raffinose, arabinose, mannose, sodium succinate, and the like.

The sources of nitrogen are yeast extract, meat extract, peptone, gluten meal, cottonseed meal, soybean meal and other vegetable meals (partially or totally defatted), casein hydrolysates, soybean hydrolysates, and yeast hydrolysates, corn steep liquor, dried yeast, wheat germ, feather meal, peanut powder, distiller's solubles, etc., as well as inorganic and organic nitrogen compounds such as ammonium salts (e.g., ammonium nitrate, ammonium sulfate, ammonium phosphate, etc.), urea, amino acids, and the like.

The carbon and nitrogen sources, though advantageously employed in combination, need not be used in their pure form, because less pure materials which contain traces of growth factors and considerable quantities of mineral nutrients, are also suitable for use. When desired, there may be added to the medium mineral salts such as sodium or calcium carbonate, sodium or potassium phosphate, sodium or potassium chloride, sodium or potassium iodide, magnesium salts, copper salts, cobalt salts, and the like. If necessary, especially when the culture medium foams seriously, a defoaming agent, such as liquid paraffin, fatty oil, plant oil, mineral oil or silicone may be added

Agitation and aeration of the culture mixture may be accomplished in a variety of ways. Agitation may be provided by a propeller or similar mechanical agitation equipment, by revolving or shaking the fermentor, by various pumping equipment, or by the passage of sterile air through the medium. Aeration may be effected by passing sterile air through the fermentation mixture

The fermentation is usually conducted at a temperature between about 20°C and 30°C, preferably 22-25°C, for a period of about 14-21 days, which may be varied according to fermentation conditions and scales.

Preferred culturing/production media for carrying out the fermentation
5 those set forth in the Examples.

As to the conditions for the production of cells in massive amounts, submerged aerobic cultural conditions is one method of culturing the cells. For the production in small amounts, a shaking or surface culture in a flask or bottle is employed. The use of fermentors (tanks) is preferred for the generation of large
10 quantities of materials. Fermentors can be sterilized with the production medium or can be sterilized empty and the medium sent through a continuous sterilizer, which is preferred for very large fermentations (20,000 gallons or larger). Preferably, the pH of the medium is adjusted to about 6-7, generally using acid or base additions, preferably made automatically with a pH electrode and a controller. The parameters
15 for fermenter operation include agitation, aeration, temperature and pressure. Agitation is preferably carried out by mixing the medium with a number of impellers mounted on a rotating agitator shaft located in the midst of the tank. Aeration may be carried out by a variety of means, preferably by bubbling sterile air into the medium (subsurface sparging). The tank is preferably maintained under positive pressure.
20 Temperature is preferably maintained at between about 20°C and 30°C.

When the growth is carried out in large tanks, vegetative forms of the organism for inoculation in the production tanks may be employed in order to avoid growth lag in the process of production. This requires production of a vegetative inoculum of the organism by inoculating a relatively small quantity of culture medium
25 with spores or mycelia of the organism produced in a "slant" and culturing said inoculated medium, also called the "seed medium", and then transferring the cultured vegetative inoculum aseptically to large tanks. The fermentation medium, in which the inoculum is produced, is generally sterilized prior to inoculation. The pH of the medium is generally adjusted to about 6-7 prior to the autoclaving step, generally
30 using acid or base additions, preferably made automatically with a pH electrode and a controller.

Preferred culturing/production media for carrying out the fermentation are those set forth in the Examples.

After growth is completed, the cells harvested by adding the
35 appropriate solvent, e.g., methylethylketone, to the entire culture medium and cells. If

the culture is grown in a liquid fermentation, the growth could be harvested by other conventional methods, e.g., centrifugation and filtration, and then extracted with the appropriate solvent, e.g., methylethylketone.

The product of the present invention can be recovered from the culture medium by conventional means which are commonly used for the recovery of other known substances. The substances produced may be found in either or both the cultured mycelium and broth filtrate, and accordingly can be isolated and purified from the mycelium and the filtrate, which are obtained by filtering or centrifuging the cultured broth, by a conventional method such as concentration under reduced pressure, lyophilization, extraction with a conventional solvent, such as methylene chloride or methanol and the like, pH adjustment, treatment with a conventional resin (e.g., anion or cation exchange resin, non-ionic adsorption resin, etc.), treatment with a conventional adsorbent (e.g., activated charcoal, silicic acid, silica gel, cellulose, alumina, etc.), crystallization, recrystallization, and the like. A preferred method is extraction of cultured whole broth with methylethylketone, followed by filtration of the extract through filtering aid such as diatomaceous earth. The methylethylketone layer of the filtrate is separated and concentrated to dryness initially by evaporating under reduced pressure followed by lyophilization. The compounds are finally isolated either by solvent partitioning, crystallization, gel filtration or by preparative HPLC on reversed phase systems.

Compounds of Formula (I) may be isolated from the aerobic fermentation of a culture of MF6388 (ATCC 74478). A culture of MF6388 (ATCC 74478) is defined as substantially free of its natural soil contaminants and capable of forming compounds of structural Formula (I) in recoverable amounts. The culture employed in the present invention should be free from viable contaminating microorganisms deleterious to the production of the compound of structural Formula (I). A biologically pure culture of MF6388 (ATCC 74478) may also be employed.

Compounds of Formula (I) may be isolated from the aerobic fermentation of a culture of MF6591 (ATCC 74477). A culture of MF6591 (ATCC 74477) is defined as substantially free of its natural soil contaminants and capable of forming compounds of structural Formula (I) in recoverable amounts. The culture employed in the present invention should be free from viable contaminating microorganisms deleterious to the production of the compound of structural Formula (I). A biologically pure culture of MF6591 (ATCC 74477) may also be employed.

The compounds of the present invention may be administered in the form of pharmaceutically acceptable salts. The term "pharmaceutically acceptable salt" is intended to include all acceptable salts such as acetate, lactobionate, benzenesulfonate, laurate, benzoate, malate, bicarbonate, maleate, bisulfate, mandelate, bitartrate, mesylate, borate, methylbromide, bromide, methylnitrate, calcium edetate, methylsulfate, camsylate, mucate, carbonate, napsylate, chloride, nitrate, clavulanate, N-methylglucamine, citrate, ammonium salt, dihydrochloride, oleate, edetate, oxalate, edisylate, pamoate (embonate), estolate, palmitate, esylate, pantothenate, fumarate, phosphate/diphosphate, gluceptate, polygalacturonate, gluconate, salicylate, glutamate, stearate, glycolylarsanilate, sulfate, hexylresorcinolate, subacetate, hydrabamine, succinate, hydrobromide, tannate, hydrochloride, tartrate, hydroxynaphthoate, teoclate, iodide, tosylate, isothionate, triethiodide, lactate, pantoate, valerate, and the like which can be used as a dosage form for modifying the solubility or hydrolysis characteristics or can be used in sustained release or pro-drug formulations. Depending on the particular functionality of the compound of the present invention, pharmaceutically acceptable salts of the compounds of this invention include those formed from cations such as sodium, potassium, aluminum, calcium, lithium, magnesium, zinc, and from bases such as ammonia, ethylenediamine, N-methyl-glutamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chlorprocaine, diethanolamine, procaine, N-benzylphenethylamine, diethylamine, piperazine, tris(hydroxymethyl)aminomethane, and tetramethylammonium hydroxide. These salts may be prepared by standard procedures, e.g. by reacting a free acid with a suitable organic or inorganic base. Where a basic group is present, such as amino, an acidic salt, i.e. hydrochloride, hydrobromide, acetate, pamoate, and the like, can be used as the dosage form.

Also, in the case of an acid (-COOH) or alcohol group being present, pharmaceutically acceptable esters can be employed, e.g. acetate, maleate, pivaloyloxymethyl, and the like, and those esters known in the art for modifying solubility or hydrolysis characteristics for use as sustained release or prodrug formulations.

For these purposes, the compounds of the present invention may be administered orally, parenterally (including subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques), by inhalation spray, or rectally, in dosage unit formulations containing conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles.

The terms "administration of" and or "administering a" compound should be understood to mean providing a compound of the invention or a prodrug of a compound of the invention to the individual in need of treatment.

Thus, in accordance with the present invention there is further
5 provided a method of treating and a pharmaceutical composition for treating HIV infection and AIDS. The treatment involves administering to a patient in need of such treatment a pharmaceutical composition comprising a pharmaceutical carrier and a therapeutically-effective amount of a compound of the present invention.

As used herein, the term "composition" is intended to encompass a
10 product comprising the specified ingredients in the specified amounts, as well as any product which results directly, or indirectly, from combination of the specified ingredients in the specified amounts.

By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not
15 deleterious to the recipient thereof.

These pharmaceutical compositions may be in the form of orally-administrable suspensions or tablets, nasal sprays, sterile injectible preparations, for example, as sterile injectible aqueous or oleagenous suspensions or suppositories.

When administered orally as a suspension, these compositions are
20 prepared according to techniques well-known in the art of pharmaceutical formulation and may contain microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners/flavoring agents known in the art. As immediate release tablets, these compositions may contain microcrystalline cellulose, dicalcium phosphate, starch,
25 magnesium stearate and lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants known in the art.

When administered by nasal aerosol or inhalation, these compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or
30 other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

The injectible solutions or suspensions may be formulated according to known art, using suitable non-toxic, parenterally-acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution or isotonic sodium chloride
35 solution, or suitable dispersing or wetting and suspending agents, such as sterile,

bland, fixed oils, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.

When rectally administered in the form of suppositories, these compositions may be prepared by mixing the drug with a suitable non-initiating excipient, such as cocoa butter, synthetic glyceride esters of polyethylene glycols, which are solid at ordinary temperatures, but liquefy and/or dissolve in the rectal cavity to release the drug.

The compounds of this invention can be administered orally to humans in a dosage range of 1 to 1000 mg/kg body weight in divided doses. One preferred dosage range is 0.1 to 200 mg/kg body weight orally in divided doses. Another preferred dosage range is 0.5 to 100 mg/kg body weight orally in divided doses. For oral administration, the compositions are preferably provided in the form of tablets containing 1.0 to 1000 milligrams of the active ingredient, particularly 1.0, 5.0, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 100.0, 150.0, 200.0, 250.0, 300.0, 400.0, 500.0, 600.0, 750.0, 800.0, 900.0, and 1000.0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

The present invention is also directed to combinations of the HIV integrase inhibitor compounds with one or more agents useful in the treatment of AIDS. For example, the compounds of this invention may be effectively administered, whether at periods of pre-exposure and/or post-exposure, in combination with effective amounts of the AIDS antivirals, immunomodulators, anti-infectives, or vaccines, such as those in the following table.

30

ANTIVIRALS

Drug NameManufacturerIndication

097	Hoechst/Bayer	HIV infection, AIDS, ARC (non-nucleoside reverse transcriptase inhibitor)
Amprenavir 141 W94 GW 141	Glaxo Wellcome	HIV infection, AIDS, ARC (protease inhibitor)
Abacavir GW 1592 1592U89	Glaxo Welcome	HIV infection, AIDS, ARC (reverse transcriptase inhibitor)
Acemannan	Carrington Labs (Irving, TX)	ARC
Acyclovir	Burroughs Wellcome	HIV infection, AIDS, ARC, in combination with AZT
AD-439	Tanox Biosystems	HIV infection, AIDS, ARC
AD-519	Tanox Biosystems	HIV infection, AIDS, ARC
Adefovir dipivoxil AL-721	Gilead Sciences Ethigen (Los Angeles, CA)	HIV infection ARC, PGL HIV positive, AIDS
Alpha Interferon	Glaxo Wellcome	Kaposi's sarcoma, HIV in combination w/Retrovir
Ansamycin LM 427	Adria Laboratories (Dublin, OH) Erbamont (Stamford, CT)	ARC

Antibody which neutralizes pH labile alpha aberrant Interferon AR177	Advanced Biotherapy Concepts (Rockville, MD)	AIDS, ARC
beta-fluoro-ddA BMS-232623 (CGP-73547)	Aronex Pharm	HIV infection, AIDS, ARC
BMS-234475 (CGP-61755)	Nat'l Cancer Institute Bristol-Myers Squibb/ Novartis	AIDS-associated diseases HIV infection, AIDS, ARC (protease inhibitor)
CI-1012 Cidofovir	Bristol-Myers Squibb/ Novartis	HIV infection, AIDS, ARC (protease inhibitor)
Curdlan sulfate Cytomegalovirus immune globin Cytovene Ganciclovir	Warner-Lambert Gilead Science	HIV-1 infection CMV retinitis, herpes, papillomavirus
	AJI Pharma USA MedImmune	HIV infection CMV retinitis
Delaviridine	Syntex	sight threatening CMV peripheral CMV retinitis
Dextran Sulfate	Pharmacia-Upjohn	HIV infection, AIDS, ARC (protease inhibitor)
ddC	Ueno Fine Chem. Ind. Ltd. (Osaka, Japan)	AIDS, ARC, HIV positive asymptomatic
Dideoxycytidine	Hoffman-La Roche	HIV infection, AIDS, ARC
ddI	Bristol-Myers Squibb	HIV infection, AIDS, ARC; combination with AZT/d4T
Dideoxyinosine		

DMP-450	AVID (Camden, NJ)	HIV infection, AIDS, ARC (protease inhibitor)
EL10	Elan Corp, PLC (Gainesville, GA)	HIV infection
Efavirenz (DMP 266) (-) 6-Chloro-4(S)- cyclopropylethynyl- 4(S)-trifluoro-methyl- 1,4-dihydro-2H-3,1- benzoxazin-2-one,	DuPont (SUSTIVA®), Merck (STOCRIN®)	HIV infection, AIDS, ARC (non-nucleoside RT inhibitor)
Famciclovir	Smith Kline	herpes zoster, herpes simplex
FTC	Emory University	HIV infection, AIDS, ARC (reverse transcriptase inhibitor)
GS 840	Gilead	HIV infection, AIDS, ARC (reverse transcriptase inhibitor)
HBV097	Hoechst Marion Roussel	HIV infection, AIDS, ARC (non-nucleoside reverse transcriptase inhibitor)
Hypericin	VIMRx Pharm.	HIV infection, AIDS, ARC
Recombinant Human Interferon Beta Interferon alfa-n3	Triton Biosciences (Alameda, CA) Interferon Sciences	AIDS, Kaposi's sarcoma, ARC ARC, AIDS

Indinavir	Merck	HIV infection, AIDS, ARC, asymptomatic HIV positive, also in combination with AZT/ddI/ddC
Compound A	Merck	HIV infection, AIDS, ARC, asymptomatic HIV positive
ISIS 2922	ISIS Pharmaceuticals	CMV retinitis
KNI-272	Nat'l Cancer Institute	HIV-assoc. diseases
Lamivudine, 3TC	Glaxo Wellcome	HIV infection, AIDS, ARC (reverse transcriptase inhibitor); also with AZT
Lobucavir	Bristol-Myers Squibb	CMV infection
Nelfinavir	Agouron Pharmaceuticals	HIV infection, AIDS, ARC (protease inhibitor)
Nevirapine	Boehringer Ingelheim	HIV infection, AIDS, ARC (protease inhibitor)
Novapren	Novaferon Labs, Inc. (Akron, OH)	HIV inhibitor
Peptide T	Peninsula Labs	AIDS
Octapeptide Sequence	(Belmont, CA)	
Trisodium Phosphonoformate	Astra Pharm. Products, Inc	CMV retinitis, HIV infection, other CMV infections
PNU-140690	Pharmacia Upjohn	HIV infection, AIDS, ARC (protease inhibitor)
Probucol	Vyrex	HIV infection, AIDS

RBC-CD4	Sheffield Med. Tech (Houston TX)	HIV infection, AIDS, ARC
Ritonavir	Abbott	HIV infection, AIDS, ARC (protease inhibitor)
Saquinavir	Hoffmann-LaRoche	HIV infection, AIDS, ARC (protease inhibitor)
Stavudine; d4T Didehydrodeoxy- thymidine	Bristol-Myers Squibb	HIV infection, AIDS, ARC
Valaciclovir	Glaxo Wellcome	genital HSV & CMV infections
Virazole	Viratek/ICN	asymptomatic HIV
Ribavirin	(Costa Mesa, CA)	positive, LAS, ARC
VX-478	Vertex	HIV infection, AIDS, ARC
Zalcitabine	Hoffmann-La Roche	HIV infection, AIDS, ARC, with AZT
Zidovudine; AZT	Glaxo Wellcome	HIV infection, AIDS, ARC, Kaposi's sarcoma, in combination with other therapies
ABT-378	Abbott	HIV infection, AIDS, ARC (protease inhibitor)
JE2147/AG1776	Agouron	HIV infection, AIDS, ARC (protease inhibitor)
T-20	Trimeris	HIV infection, AIDS, ARC (fusion inhibitor)
T-1249		
BMS 232632	Bristol-Myers-Squibb	HIV infection, AIDS, ARC (protease inhibitor)

IMMUNO-MODULATORS

<u>Drug Name</u>	<u>Manufacturer</u>	<u>Indication</u>
AS-101	Wyeth-Ayerst	AIDS
Bropiramine	Pharmacia Upjohn	advanced AIDS
Acemannan	Carrington Labs, Inc. (Irving, TX)	AIDS, ARC
CL246,738	American Cyanamid Lederle Labs	AIDS, Kaposi's sarcoma
EL10	Elan Corp, PLC (Gainesville, GA)	HIV infection
FP-21399	Fuki ImmunoPharm	blocks HIV fusion with CD4+ cells
Gamma Interferon	Genentech	ARC, in combination w/TNF (tumor necrosis factor)
Granulocyte Macrophage Colony Stimulating Factor	Genetics Institute Sandoz	AIDS
Granulocyte Macrophage Colony Stimulating Factor	Hoeschst-Roussel Immunex	AIDS
Granulocyte Macrophage Colony Stimulating Factor	Schering-Plough	AIDS, combination w/AZT
HIV Core Particle Immunostimulant	Rorer	seropositive HIV
IL-2	Cetus	AIDS, in combination w/AZT
Interleukin-2		
IL-2	Hoffman-La Roche	AIDS, ARC, HIV, in combination w/AZT
Interleukin-2	Immunex	
IL-2	Chiron	AIDS, increase in CD4 cell counts
Interleukin-2 (aldeslukin)		

Immune Globulin Intravenous (human)	Cutter Biological (Berkeley, CA)	pediatric AIDS, in combination w/AZT
IMREG-1	Imreg (New Orleans, LA)	AIDS, Kaposi's sarcoma, ARC, PGL
IMREG-2	Imreg (New Orleans, LA)	AIDS, Kaposi's sarcoma, ARC, PGL
Imuthiol Diethyl Dithio Carbamate	Merieux Institute	AIDS, ARC
Alpha-2 Interferon	Schering Plough	Kaposi's sarcoma w/AZT, AIDS
Methionine- Enkephalin	TNI Pharmaceutical (Chicago, IL)	AIDS, ARC
MTP-PE	Ciba-Geigy Corp.	Kaposi's sarcoma
Muramyl-Tripeptide Granulocyte Colony Stimulating Factor	Amgen	AIDS, in combination w/AZT
Remune rCD4 Recombinant Soluble Human CD4 rCD4-IgG hybrids	Immune Response Corp. Genentech	immunotherapeutic AIDS, ARC
Recombinant Soluble Human CD4 Interferon Alfa 2a	Biogen Hoffman-La Roche	AIDS, ARC Kaposi's sarcoma AIDS, ARC, in combination w/AZT
SK&F106528 Soluble T4 Thymopentin	Smith Kline Immunobiology Research Institute	HIV infection HIV infection

Tumor Necrosis Factor; TNF etanercept infliximab	Genentech Immunex Corp (Enbrel®) Centocor (Remicade®)	ARC, in combination w/gamma Interferon rheumatoid arthritis rheumatoid arthritis and Crohn's disease
---	---	--

ANTI-INFECTIVES

<u>Drug Name</u>	<u>Manufacturer</u>	<u>Indication</u>
Clindamycin with Primaquine Fluconazole	Pharmacia Upjohn Pfizer	PCP cryptococcal meningitis, candidiasis
Pastille Nystatin Pastille Ornidyl Eflornithine	Squibb Corp. Merrell Dow	prevention of oral candidiasis PCP
Pentamidine Isethionate (IM & IV) Trimethoprim Trimethoprim/sulfa Piritrexim Pentamidine isethionate for inhalation Spiramycin	LyphoMed (Rosemont, IL) Burroughs Wellcome Fisons Corporation Rhone-Poulenc	PCP treatment antibacterial antibacterial PCP treatment PCP prophylaxis cryptosporidial diarrhea
Intraconazole- R51211 Trimetrexate	Janssen Pharm. Warner-Lambert	histoplasmosis; cryptococcal meningitis PCP

<u>Drug Name</u>	<u>Manufacturer</u>	<u>Indication</u>
Daunorubicin	NeXstar, Sequus	Kaposi's sarcoma
Recombinant Human Erythropoietin	Ortho Pharm. Corp.	severe anemia assoc. with AZT therapy
Recombinant Human Growth Hormone	Serono	AIDS-related wasting, cachexia
Leukotriene B4 Receptor Antagonist	-	HIV infection
Megestrol Acetate	Bristol-Myers Squibb	treatment of anorexia assoc. w/AIDS
Soluble CD4 Protein and Derivatives	-	HIV infection
Testosterone	Alza, Smith Kline	AIDS-related wasting
Total Enteral Nutrition	Norwich Eaton Pharmaceuticals	diarrhea and malabsorption related to AIDS

It will be understood that the scope of combinations of the compounds of this invention with HIV/AIDS antivirals, immunomodulators, anti-infectives or vaccines is not limited to the list in Table 1 above, but includes in principle any combination with any pharmaceutical composition useful for the treatment of AIDS.

One preferred combination is a compound of the present invention and a nucleoside inhibitor of HIV reverse transcriptase such as AZT, 3TC, ddC, or ddI. Another preferred combination is a compound of the present invention and a non-nucleoside inhibitor of HIV reverse transcriptase, such as efavirenz, and optionally a nucleoside inhibitor of HIV reverse transcriptase, such as AZT, 3TC, ddC or ddI. Still another preferred combination is any one of the foregoing combinations further comprising an additional HIV protease inhibitor such as indinavir, Compound A, nelfinavir, ritonavir, saquinavir, amprenavir, or abacavir. A preferred additional inhibitor of HIV protease is the sulfate salt of indinavir. Other preferred additional protease inhibitors are nelfinavir and ritonavir. Still another preferred additional inhibitor of HIV protease is saquinavir which is administered in a dosage of 600 or 1200 mg tid.

Other preferred combinations include a compound of the present invention with the following (1) efavirenz, optionally with AZT and/or 3TC and/or ddI and/or ddC, and optionally with indinavir; (2) any of AZT and/or ddI and/or ddC and/or 3TC, and optionally with indinavir; (3) d4T and 3TC and/or AZT; (4) AZT and 3TC; and (5) AZT and d4T.

In such combinations the compound of the present invention and other active agents may be administered together or separately. In addition, the administration of one agent may be prior to, concurrent to, or subsequent to the administration of other agent(s). These combinations may have unexpected effects on limiting the spread and degree of infection of HIV.

Efavirenz is (-)-6-chloro-4-cyclopropylethynyl-4-trifluoromethyl-1,4-dihydro-2H-3,1-benzoxazin-2-one, also known as DMP-266 or SUSTIVA® (DuPont) or STOCRIN® (Merck). Efavirenz and its utility as an HIV reverse transcriptase inhibitor is described in US 5519021 and in the corresponding PCT published application, WO 95/20389. Efavirenz can be synthesized by the protocol of US 5633405. Additionally, the asymmetric synthesis of an enantiomeric benzoxazinone by a highly enantioselective acetylide addition and cyclization sequence is described in Thompson et al., *Tetrahedron Letters* 1995, 36: 8937-40, as well as in the PCT publication, WO 96/37457.

AZT is 3'-azido-3'-deoxythymidine, is also known as zidovudine, and is available from Burroughs-Wellcome under the tradename RETROVIR®. Stavudine is 2',3'-didehydro-3'-deoxythymidine, is also known as 2',3'-dihydro-3'-deoxythymidine and d4T, and is available from Bristol-Myers Squibb under the tradename ZERIT®. 3TC is (2R-cis)-4-Amino-1-[2-(hydroxymethyl)-1,3-oxathiolan-5-yl]-2(1H)-pyrimidinone, is also known as (-)-1-[(2R,5S)-2-(hydroxymethyl)-1,3-oxathiolan-5-yl]cytosine and lamivudine, and is available from Glaxo Wellcome under the tradename EPIVIR®. ddC is 2',3'-dideoxycytidine, is also known as zalcitabine, and is available from Hoffman LaRoche under the tradename HIVID®. ddI is 2',3'-dideoxyinosine, is also known as didanosine, and is available from Bristol-Myers-Squibb under the tradename VIDEX®. The preparation of ddC, ddI and AZT are also described in EPO 0,484,071.

Indinavir is N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4-(S)-hydroxy-5-(1-(4-(3-pyridyl-methyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide, and can be prepared as described in US 5413999. Indinavir is

generally administered as the sulfate salt at a dosage of 800 mg three times a day. Indinavir is available from Merck under the tradename CRIVAN®.

Compound A is N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(2-benzo[b]furanylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))pentaneamide, preferably administered as the sulfate salt. Compound A can be prepared as described in US 5646148.

Ritonavir is [5S-(5R*,8R*,10R*, 11R*)]-10-hydroxy-2-methyl-5-(1-methylethyl)-1-[2-(1-methylethyl)-4-thiazolyl]-3,6-dioxo-8,11-bis(phenylmethyl)-2, 4, 7, 12-tetraazatridecan-13-oic acid 5-thiazolylmethyl ester, also known as 5-thiazolylmethyl [(aS)-a-[(1S,3S)-1-hydroxy-3-[(2S)-2-[3-[(2-isopropyl-4-thiazolyl)methyl]-3-methylureido]-3-methylbutyramido]-4-phenylbutyl]phenethyl]carbamate. It is available from Abbott under the tradename NORVIR®. Ritonavir can be prepared as described in US 5484801.

Nelfinavir is [3S-[2(2S*,3S*),3a,4ab,8ab]]-N-(1,1-dimethylethyl)decahydro-2-[2-hydroxy-3-[(3-hydroxy-2-methylbenzoyl)amino]-4-(phenylthio)butyl]-3-isoquinolinecarboxamide monomethanesulfonate, also known as (3S,4aS,8aS)-N-tert-Butyl-2-[(2R,3R)-3-(3,2-crestamido)-2-hydroxy-4-(phenylthio)butyl]decahydro-3-isoquinolinecarboxamide monomethanesulfonate and VIRACEPT®, which is commercially available from Agouron. Nelfinavir can be prepared as described in US 5484926.

Saquinavir is N-tert-butyl-decahydro-2-[2(R)-hydroxy-4-phenyl-3(S)-[[N-(2-quinolylcarbonyl)-L-asparaginy]amino]butyl]-(4aS,8aS)-isoquinoline-3(S)-carboxamide, also known as INVIRASE®. Saquinavir can be prepared in accordance with procedures disclosed in US 5196438. INVIRASE® (saquinavir mesylate) is available from Roche Laboratories. Saquinavir can be prepared as described in US 5196438.

Amprenavir is 4-amino-N-((2 syn,3S)-2-hydroxy-4-phenyl-3-((S)-tetrahydrofuran-3-ylloxycarbonylamino)-butyl)-N-isobutyl-benzenesulfonamide, also known as Compound 168 and 141 W94. Amprenavir is an aspartyl protease inhibitor that can be prepared by following the procedures described in US 5585397. Amprenavir is available under the tradename AGENERASE® from Glaxo Wellcome. Amprenavir can be prepared as described in US 5783701.

Abacavir is (1S,4R)-cis-4-[2-amino-6-(cyclopropylamino)-9H-purin-9-yl]-2-cyclopentene-1-methanol, also known as 1592U89. Abacavir can be prepared by following the protocol of EP 0434450. The following examples are

provided to further illustrate details for the preparation and use of the compounds of the present invention. The examples are not intended to be limitations on the scope of the instant invention in any way, and they should not be so construed.

Furthermore, the compounds described in the following examples are not to be construed as forming the only genus that is considered as the invention, and any combination of the compounds or their moieties may itself form a genus. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds. All temperatures are in degrees Celsius unless noted otherwise.

Abbreviations: DMF is dimethyl formamide; ESIMS represents Electron Spray Ionization Mass Spectroscopy; Et represents ethyl; HPLC is high pressure liquid chromatography; HREIMS represents High Resolution Electron Impact Mass Spectroscopy; IPA is isopropyl alcohol; MEK is methyl ethyl ketone; Me represent methyl; rh represents relative humidity, TFA is trifluoroacetic acid; THF is tetrahydrofuran; TLC is thin layer (SiO₂) chromatography.

EXAMPLE 1

Fermentation of MF6388 (ATCC 74478) *Sterile Fungus*

A. Media

SEED MEDIUM:

	<u>Component</u>	<u>g/L</u>
	Yeast extract	4.0
	Malt extract	8.0
25	Glucose	4.0
	Junlon	1.5

The medium was prepared with distilled water, the pH adjusted to 7.0 prior to sterilization, and was dispensed at 50 mL/250 mL unbaffled Erlenmeyer flask. Cotton closures were used. Sterilization was at 121°C for 20 minutes.

30 PRODUCTION MEDIUM:

1. Solid portion:

675 cc vermiculite was added to a 2 liter roller bottle which was plugged with latex closure and autoclaved for 60 minutes, plus 30 minutes and dried.

2. Liquid portion

	<u>Component</u>	<u>Amount</u>
--	------------------	---------------

	Glucose	150.0 g/L
	Fructose	15.0 g/L
	Sucrose	40.0 g/L
	Casamino acids	2.0 g/L
5	Asparagine	2.0 g/L
	Yeast extract	1.0 g/L
	Na ₂ HPO ₄	0.5 g/L
	MgSO ₄ ·7H ₂ O	1.0 g/L
	CaCl ₂	0.5 g/L
10	K-elements	1.0 mL/L
	pH to 7.0	

The medium was prepared with distilled water, dispensed at 220 mL in 500 mL bottles and sterilized at 121°C for 20 minutes.

15	K-elements	
	<u>Component</u>	<u>Amount</u>
	FeCl ₃ ·6H ₂ O	5.8 g/L
	MnSO ₄ ·H ₂ O	0.1 g/L
	CoCl ₂ ·6H ₂ O	0.02 g/L
20	CuSO ₄ ·5H ₂ O	0.015 g/L
	NaMoO ₄ ·2H ₂ O	0.012 g/L
	ZnCl ₂	0.02 g/L
	SnCl ₂ ·2H ₂ O	0.005 g/L
	H ₃ BO ₃	0.01 g/L
25	KCl	0.02 g/L
	HCl (concentrated)	2.0 mL/L

B. Inoculum Preparation

- An agar slant was used to prepare FVMs (frozen vegetative mycelia).
- 30 A portion of the agar slant was transferred aseptically to seed medium. (The composition of the seed medium is detailed above). The flasks were incubated on a 2-inch throw gyratory shaker, 220 rpm for 7 days at 25°C, 85% relative humidity (rh), to obtain biomass. Portions of the biomass were transferred into sterile vials containing glycerol and frozen (as FVM). These were maintained in a final concentration of 10-
- 35 15% glycerol at -75°C.

C. Seed Culture

Frozen vials (FVM) were thawed to room temperature and used to inoculate seed cultures, at 1.0 mL per 50 mL seed medium. The cultures were grown on a gyratory shaker (220 rpm) for 4 days at 25°C, 85% rh, until a sufficient amount
5 of biomass was obtained.

D. Production

The composition of the solid substrate fermentation medium is shown above. An aliquot (12 mL) of each grown seed was placed into 220 mL of the liquid portion of the production. This was swirled vigorously to disperse the biomass. The
10 contents were dispensed by pouring into a 2-liter roller culture vessel which contained 675 cubic centimeters of steam-sterilized large-particle vermiculite. The contents of the roller bottle were shaken/mixed to insure homogeneous inoculation and coverage. The roller bottles were incubated horizontally, revolving at approximately 4 rpm on a Wheaton roller apparatus at 22°C, 75% rh for 20 days, to obtain secondary metabolite
15 production in the fermentation medium. The contents of each roller bottle were extracted with 200-250 mL methyl ethyl ketone and the solids discarded.

EXAMPLE 2

Fermentation of MF6591 (ATCC 74477) *Ascochyta sp.*

20 A. Media:

Seed medium contained the following in g/L: corn steep liquor, 5g; tomato paste, 40; oat flour, 10; glucose, 10; agar, 4; FeSO₄·7H₂O, 0.01; MnSO₄·4H₂O, 0.01; CuCl₂·2H₂O, 0.00025; CaCl₂, 0.001; H₃BO₃, 0.00056; (NH₄)₆Mo₇O₂₄·4H₂O, 0.00019; ZnSO₄·7H₂O, 0.002. The pH was adjusted to 6.8.

25 Production media contained the following per 250 mL flask: brown rice, 10g; base liquid, 20 mL. Base liquid contained the following in g/L: yeast extract, 1; sodium tartrate, 0.5; KH₂PO₄, 0.5. The flasks were autoclaved for 15 minutes at 121°C, 15 psi and stored. Prior to inoculation, 15 mL of distilled water were added per flask and the flasks were sterilized for 20 minutes at 121° C, 15 psi.

30 B. Inoculum Preparation

Frozen vegetative mycelia (FVM) were prepared by inoculating 50 mL of seed medium in a 250 mL flask and incubating at 25°C, 85% relative humidity and at 200 rpm for 3-5 days. Aliquots of the culture were frozen and used as a source of inoculum for future experiments.

35 C. Seed Culture

To 50 mL of seed media in a 250 mL flask, 2.0 mL of FVM was added as inoculum and the flasks were incubated at 25°C, 85% relative humidity and at 200 rpm for days 2-3 days.

D. Production of Culture and Extraction

- 5 To 50 mL of production media in a 250 mL flask, 1 mL of seed culture was added as inoculum and the flasks were incubated at 25°C, 85% relative humidity for 24 days. Each flask was then extracted with 50 mL of methyl ethyl ketone and the solids were discarded.

10

EXAMPLE 3

Isolation of Compounds A and B from MF6591:

- The fungus was grown on a brown rice based liquid medium for 31 days and 1.4 L of the broth was extracted with 1.2 volumes of methyl ethyl ketone (MEK). The layers were separated to give 1.4 L MEK extract. The extract was
- 15 concentrated under reduced pressure to give an aqueous solution that was diluted by addition of 300 mL of water and 50 mL of methanol. This methanolic aqueous solution was partitioned three times with one volume each of hexane followed by five times with one volume each of ethyl acetate. All of the integrase activity was concentrated in ethyl acetate extract. This extract was concentrated to dryness to give
- 20 a brown gum. Five grams of this material was chromatographed on a 2.0 L SEPHADEX LH20 column. Elution of the column with methanol eluted the activity in a broad zone from 0.6 L to 1.7 L column volume. The most active fraction contained mostly the mixture of Compounds A and B. An aliquot of this fraction was chromatographed on a reverse phase HPLC using ZORBAX RX C-8 (22 x 250 mm)
- 25 column. The column was eluted at 8 mL/min with a gradient of acetonitrile in water containing 0.05% trifluoroacetic acid. The gradient started with 20% acetonitrile to 40% acetonitrile in 50 minutes, held at 40% for 10 minutes and then the acetonitrile concentration was increased to 100% over 20 minutes. Lyophilization of fractions eluting in between 31-35 and 52-59 minutes gave Compounds B and A, respectively,
- 30 as brown powders. Compound B: UV λ_{max} : 210, 230, 315 nm. ESIMS (m/z) 405 (M+H)⁺, 403 (M-H)⁻; HREIMS (m/z): 404.1110 (M⁺, calcd for C₂₀H₂₀O₉: 404.1107), 209.0436 (calcd for C₁₀H₉O₅: 209.0450), 195.0623 (calcd for C₁₀H₁₁O₄: 195.0657); for ¹H and ¹³C NMR see Table 1. Compound A: UV λ_{max} : 215, 235, 320 nm. ESIMS (m/z) 403 (M+H)⁺, 401 (M-H)⁻; HREIMS (m/z): 402.0960


(M⁺, calcd for C₂₀H₁₈O₉: 402.0950), 209.0452 (calcd for C₁₀H₉O₅: 209.0450), 194.0576 (calcd for C₁₀H₁₁O₄: 194.0579); for ¹H and ¹³C NMR see Table 1.

EXAMPLE 4

5 Isolation of Compound A from MF6388:

Sixty mL methyl ethyl ketone extract of MF6388 grown on vermiculite solid medium was concentrated to dryness to give a brown gum. This material was dissolved in a 1:1 mixture of methylene chloride and methanol and chromatographed on a 1.0 L column of SEPHADEX LH20. The column was eluted with methanol. A single combined active fraction was chromatographed on a ZORBAX RX C-8 reverse phase HPLC column. Gradient elution with 20 to 80% acetonitrile in water containing 0.05% TFA over 40 minutes at a flow rate of 8 mL/min gave after lyophilization Compound A as a brown powder.

15 Table 1: ¹H NMR and ¹³C NMR spectral data of Compounds B and A



Compound B; R = CH₂OH

Compound A; R = CHO

Position	δ C Compound B CD ₃ CN	δ H Compound B CD ₃ CN	δ C Compound A CDCl ₃ + CD ₃ CN	δ H Compound A CDCl ₃ + CD ₃ CN
2	97.8		97.8	
3	121.3		120.7	
4	140.1*		140.1*	
5	140.4		142.2	
6	148.6		148.7	

7	101.4	7.03, s	101.8	7.06, s
8	121.1		120.4	
9	193.8		193.7	
10	77.4		77.1	
11	114.1		121.3	
12	130.5		126.0	
13	107.3	6.70, s	105.7	7.14, s
14	147.7		147.8	
15	134.4		140.9	
16	142.0*		140.8*	
18	26.4	2.09, s	26.5	2.15, s
19	22.2	1.76, s	25.8	1.87, s
20	60.9	4.58, d, 12.8 4.40, d, 12.8	190.6	10.21, s
21	56.4	3.75, s	56.6	3.75, s
22	56.6	3.83, s	56.7	3.85, s

EXAMPLE 5

HIV Integrase Assay: Strand Transfer Catalyzed by Recombinant Integrase

5 Assays for the strand transfer activity of integrase were conducted according to Wolfe, A.L. et al., J. Virol. 70, 1424 (1996), hereby incorporated by reference for these purposes. Representative compounds tested in the integrase assay demonstrated IC₅₀'s less than 10 micromolar.

EXAMPLE 6

Oral Composition


10 As a specific embodiment of an oral composition of a compound of this invention, 50 mg of a compound of the present invention is formatted with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a
15 size 0 hard gelatin capsule.

 While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be

understood that the practice of the invention encompasses all of the usual variations, adoptions, or modifications, as come within the scope of the following claims and their equivalents.

WHAT IS CLAIMED:

1. A compound of Formula I:




wherein:

R is selected from:

- (a) $-\text{CH}_2\text{OH}$, and
 (b) $-\text{CH}(=\text{O})$;


or a pharmaceutically acceptable salt thereof.

2. The compound according to Claim 1, which is:



or a pharmaceutically acceptable salt thereof.

3. The compound according to Claim 1, which is:



or a pharmaceutically acceptable salt thereof.

4. A pharmaceutical composition comprising an effective amount of a compound according to Claim 1, or a pharmaceutically acceptable salt thereof,
5 and a pharmaceutically acceptable carrier.

5. A pharmaceutical composition comprising a therapeutically effective amount of a compound of Claim 1, or a pharmaceutically acceptable salt thereof, in combination with a therapeutically effective amount of an AIDS treatment
10 agent selected from:

- (a) an AIDS antiviral agent,
- (b) an immunomodulator, and
- (c) an anti-infective agent.

15 6. The pharmaceutical composition according to Claim 5 wherein the AIDS antiviral agent is N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyll))-pentaneamide, or a pharmaceutically acceptable salt thereof.

20 7. A pharmaceutical composition made by combining the compound of Claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

25 8. A process for making a pharmaceutical composition comprising combining a compound of Claim 1, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

9. A method of inhibiting HIV integrase, comprising the administration to a mammal in need of such treatment a therapeutically effective
30 amount of a compound of Claim 1, or a pharmaceutically acceptable salt thereof.

10. A method of treating infection by HIV, or of treating AIDS or ARC, comprising the administration to a mammal in need of such treatment a therapeutically effective amount of a compound of Claim 1, or a pharmaceutically
35 acceptable salt thereof.

11. The method according to Claim 10 additionally comprising the administration of a therapeutically effective amount of another AIDS treatment agent selected from:

- 5 (a) an AIDS antiviral agent,
 (b) an immunomodulator, and
 (c) an anti-infective agent.


12. The method according to Claim 11 wherein the AIDS antiviral agent is N-(2(R)-hydroxy-1(S)-indanyl)-2(R)-phenylmethyl-4(S)-hydroxy-5-(1-(4-(3-pyridylmethyl)-2(S)-N'-(t-butylcarboxamido)-piperazinyl))-pentaneamide, or a pharmaceutically acceptable salt thereof.

13. A biologically pure culture of MF6388 (ATCC 74478).

15

14. A culture of MF6388 (ATCC 74478).

15. A biologically pure culture of Claim 13, or a mutant thereof, capable of producing a compound of structural Formula (I)



20

(I)


wherein:

R is selected from:

- 25 (a) -CH₂OH, and
 (b) -CH(=O);

in recoverable amounts.

16. A culture of Claim 14, or a mutant thereof, capable of producing a compound of structural Formula (I)



(I)

wherein:

5

R is selected from:

- (a) $-\text{CH}_2\text{OH}$, and
- (b) $-\text{CH}(=\text{O})$;

in recoverable amounts.

10


17. A process of making a compound according to Claim 1 comprising cultivating MF6388 (ATCC 74478) or a mutant thereof under conditions suitable for formation of the compound and recovering the compound.

15

18. A biologically pure culture of MF6591 (ATCC 74477).

19. A culture of MF6591 (ATCC 74477).

20. A biologically pure culture of Claim 18, or a mutant thereof, capable of producing a compound of structural Formula (I)



(I)


wherein:

R is selected from:

20

- (a) $-\text{CH}_2\text{OH}$, and
 (b) $-\text{CH}(=\text{O})$;
 in recoverable amounts.

- 5 21. A culture of Claim 19, or a mutant thereof, capable of producing a compound of structural Formula (I)



(I)

- 10 wherein:

R is selected from:

- (a) $-\text{CH}_2\text{OH}$, and
 (b) $-\text{CH}(=\text{O})$;
 in recoverable amounts.

- 15 22. A process of making a compound according to Claim 1 comprising cultivating MF6591 (ATCC 74477) or a mutant thereof under conditions suitable for formation of the compound and recovering the compound.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/20918

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : C07D 325/00; C12P 17/18; A61K 31/335 US CL : 549/354; 514/450; 435/119 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 549/354; 514/450; 435/119 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) stn		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,726,203 A (LI et al.) 10 March 1998, column 3, lines 41-59.	1-22
A	US 5,616,609 A (IKEKAWA et al.) 01 April 1997, column 1, line 40 thru column 2, line 13.	1-22
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *Z* document member of the same patent family	
Date of the actual completion of the international search 31 OCTOBER 2000		Date of mailing of the international search report 30 NOV 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer AMELIA A. OWENS Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/20918**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐

The additional search fees were accompanied by the applicant's protest.

☒

No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/20918

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claim(s) 1-12, drawn to compounds of formula I, compositions, method of preparing, method of inhibiting hiv integrase, method of treating infection by hiv.

Group II, claim(s) 13-17, drawn to culture MF6388 and compound produced thereby.

Group III, claim(s) 18-22 drawn to culture MF6591 and compound produced thereby..

The inventions listed as Groups I - III do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The cultures of groups II and III of may produce other metabolites than those of group I.